

Thermodynamics of the Bose-System with a Small Number of Particles

I. O. Vakarchuk*, A. A. Rovenchak†

Department for Theoretical Physics of Lviv National University,
12 Draghomanov St., UA-79005, Lviv, Ukraine

A theoretical description of the interacting Bose-system is proposed. It is based on the extrapolation of the results obtained for the systems with a small number of particles $N = 2, 3, 4$, etc. to the bulk case of $N = \infty$. It is shown that already the system with $N = 12, 13$ behaves almost as a bulk in a wide temperature range. Special attention is paid to the phase transition in these systems. The hard sphere potential is used in calculations. The sequence of heat capacity maxima is approximated as $C_N^{\max}/N \simeq 13.6 - aN^{-\varepsilon}$ with $\varepsilon = 0.0608$ giving the value of bulk heat capacity as 13.6 while experimental value is close to 16. The temperature of λ -transition is estimated as 2.1–2.3 K versus experimental 2.17 K. Quite good qualitative and satisfactory quantitative agreement with the experimental data has been achieved.

Keywords: Boson systems, finite systems, lambda-transition, specific heat.

I. INTRODUCTION

Liquid ^4He is the only atomic system having its properties determined mainly by quantum effects and the most essential among the latter is the statistical effect of identical particles. Due to it the macroscopic number of the ^4He atoms can have a zero momentum at some conditions, i. e., it can occupy a single quantum state. The phenomenon of bosons accumulation at the lowest energy level is called Bose-Einstein condensation (1924). According to London's assumption (1938), it is believed that superfluidity, λ -transition, and other unusual properties of liquid ^4He are a consequence of this very effect.

However, despite the lapse of time that has passed since the discovery the phenomena of λ -transition by Keesom and Clausius in 1932 and the superfluidity by Kapitsa in 1938 until now no one has managed to create a comprehensive microscopic theory explaining the properties of strongly interacting Bose-systems such as ^4He within the whole temperature range of $0 < T < \infty$.

Bogoliubov [1] was the first to show how one can build a theory of real helium proceeding from the approximate second quantization techniques for the model of weakly-imperfect Bose-gas. This theory works well for $T \lesssim 1$ K, especially while $T \rightarrow 0$.

The subsequent theoretical studies brought about the appearance of the quantitative theory of the ground state of a strongly-imperfect many-boson system leading to a good agreement with the experiment and the creation of the theoretical description of low-energy excitations which are responsible for the low-temperature behaviour of thermodynamic functions. The experimental investigations of the spatial structure by means of X-ray and neutron diffraction and of the structure of the energy spectrum and the condensate fraction by means of the inelastic neutron scattering revealed a fairly good theoretical explanation.

As a consequence of the experimental evidences it was supposed for a long time that ^4He heat capacity had a logarithmic divergency at the transition temperature T_λ . Brout showed in his early works [2] based on the Hartree-Fock method that it was impossible to find an explanation of the observed heat capacity behaviour within simple theoretical approaches. There is an interesting notice regarding this question Feynman made in his book [3]: "The explanation of this behavior is left as an exercise for the reader. If successful, publish!"

Feynman suggested that the λ -transition should be studied using the full N -particle density matrix for the ideal Bose-gas with a phenomenologically introduced effective mass for particles [4]. This approach leads only to a shift of the Bose-condensation temperature in comparison with the ideal case leaving the form of the heat capacity curve unchanged.

In the early 1970s the studies of the λ -transition were started in the renormalization group (RG) approach for the theory with the degenerate two-component order parameter. The rigorous grounds of this approach based on the

*chair@ktf.franko.lviv.ua

†andrij@ktf.franko.lviv.ua

functional integration techniques are given in Refs. 5, 6. They use the coherent states representation with the complex order parameter the absolute value of which equals $\sqrt{N_0}$ where N_0 is the number of atoms in Bose-condensate.

In this approach the heat capacity critical exponent was calculated as $\alpha = 1/8$. It obviously does not correspond to the logarithmic divergence for which α equals zero. However, taking into account that the calculated value of α is a small magnitude, the experimental confirmation of this result required very precise measurements which became available only recently [7]. Before it, there was even no certainty about the sign of α . The measurements showed that the critical exponent α is negative, i. e., the heat capacity is finite at the transition temperature.

Recently the theoretical confirmation of this result appeared. Kleinert [8,9] obtained for α the respective values of $\alpha = -0.01294 \pm 0.00060$ and $\alpha = -0.0120 \pm 0.0009$, while the experiment gives $\alpha = -0.01285 \pm 0.00038$ [7]. One can find a bit different approach to the critical exponents calculation in the paper by Campostrini *et al* [10]. The authors obtained the value of $\alpha = -0.0150 \pm 0.0017$. But, despite a very good agreement of these results with the experiment, RG works only in the immediate vicinity of the transition point T_λ and gives no possibility of getting some information of the thermodynamic functions behaviour in a wider near-critical range, e. g., for $|T - T_\lambda| \sim 10^{-3}$ K.

In addition to these theoretical investigations, the numerical studies of ^4He thermodynamic properties were conducted. The Diffusion Monte Carlo (DMC) techniques are applied for the ground state calculations, and the Path Integral Monte Carlo (PIMC) is used for finite temperatures. These techniques have been applied since the 1960s. At present, PIMC is the most effective method. Its only essential imperfection lies in the fact that it is impossible to use it for the ground state ($T = 0$ K). But one can avoid this problem by considering the limit of the results at $T \rightarrow 0$ K. A detailed analysis of PIMC might be found in the work by Ceperley [11]. It appears that the numerical approach provides a good agreement with the experimental data in a wide temperature range on the whole, except for the vicinity of the λ -transition point.

Thus, we have the situation when no satisfactory theoretical description of ^4He exists for a considerable temperature range of $1 \text{ K} \lesssim T < T_\lambda$ and $T > T_\lambda$, and the only way out is the application of numerical techniques. It gives some space for the investigations in this domain. Not claiming the solution of the problem, we still propose here the model which, in our opinion, can near us to the understanding of the processes in many-boson system via the consideration of systems with a small number of particles.

For the latter one can expect a much more precise description in terms of statistical mechanics in comparison with what might be reached while directly considering a many-particle system.

The aim of the present work consists in obtaining a thermodynamic description of an interacting many-boson system basing on the results for the systems with a small number of particles N . We start our analysis from $N = 2, 3, 4$, etc. in order to extrapolate these results for the case of $N = \infty$ after establishing some dependencies on N and the temperature. We pay a special attention to the fact that even the system consisting of a small number of particles exhibits the peculiarities on the heat capacity curve such as the maximum which means the phase transition in the limit of $N \rightarrow \infty$. We will show at the same time that the system with $N = 12, 13$ tends to reveal qualitatively an “almost bulk” behaviour in a wide temperature range.

The boson systems with a relatively small number of particles have been experimentally obtained only in the last couple of years as particles in traps (see Refs. 12, 13 and references therein). It gives us a possibility of developing some methods for such systems and a further comparison of the results with the experimental data.

We use the partition function formalism to calculate thermodynamic characteristics of the free Bose-system. A method somewhat similar to this one was used in works [12,13] for the study of the trapped finite Bose-systems: the recursion formulae for the partition function proposed by P. Borrmann and G. Franke [14]. But our expressions do not take the restriction for the system energy to be written as a sum of one-particle energies. So, one can use them not only for the ideal gas but for the interacting system as well.

The paper is organized as follows. In Section II some known initiating expressions for the calculating the thermodynamic function are given in short. In Sections III and IV we consider how the proposed method works in the case of an ideal Bose-gas the results for which are well-known in the thermodynamic limit when the number of particles $N \rightarrow \infty$. The next step is the determination of the asymptotic behaviour of our results depending on the number of particles N . We will try to expand the found regularities for the case of interacting bosons. The corresponding calculations for the hard-sphere potential are given in Section V.

Generally, it is impossible to obtain a non-analyticity point of the heat capacity function having finite number of particles. Therefore, no “pure” phase transition can be detected in this model. But we expect a certain qualitative agreement with the experiment to be quite sufficient for the approximation used.

II. GENERAL STATEMENTS

We consider the system of N particles with the Hamiltonian $\hat{H}_N$. The partition function $Z_N = \text{Sp } e^{-\beta \hat{H}_N}$ where β is inverse temperature, $\beta = 1/T$.

In the coordinate representation we have

$$Z_N = \int d\mathbf{r}_1 \dots \int d\mathbf{r}_N R_N(\mathbf{r}_1, \dots, \mathbf{r}_N | \mathbf{r}_1, \dots, \mathbf{r}_N), \quad (1)$$

where R_N is the density matrix. The particles are bosons of the mass m (^4He atoms) with their coordinates $\mathbf{r}_i$ restricted in volume V . In this case R_N is given by [15]

$$R_N(\mathbf{r}_1, \dots, \mathbf{r}_N | \mathbf{r}'_1, \dots, \mathbf{r}'_N) = P_N(\mathbf{r}_1, \dots, \mathbf{r}_N) \frac{1}{N!} \frac{1}{\lambda^{3N}} \sum_Q \exp \left(-\frac{\pi}{\lambda^2} \sum_{j=1}^N (\mathbf{r}_j - \mathbf{r}'_{Q_j})^2 \right), \quad (2)$$

where Q is counting permutations of $(1, \dots, N)$ and λ is the thermal de Broglie wavelength

$$\lambda = \left(\frac{2\pi\beta\hbar^2}{m} \right)^{1/2}. \quad (3)$$

The factor P_N takes into account the interaction and will be described further.

The free energy F_N is given by

$$F_N = -\frac{1}{\beta} \ln Z_N, \quad (4)$$

total energy

$$E_N = \frac{\partial}{\partial \beta} (\beta F_N), \quad (5)$$

and heat capacity

$$C_N = \beta^2 \left[\frac{Z_N''}{Z_N} - \left(\frac{Z_N'}{Z_N} \right)^2 \right], \quad (6)$$

the primes mean derivation over β .

III. PARTITION FUNCTION FOR THE SYSTEM OF FREE PARTICLES

The ideal Bose-gas is a well-studied system for which the expressions for the thermodynamic functions are known. Thus, it can be a good test example for the applicability of our method and we are going to use this fact here.

On calculating the thermodynamic functions of the ideal Bose-system with a small number of particles and comparing them with the well-known results for infinite system one can detect some regularities showing the point when several particles start to demonstrate the statistical behaviour.

One can find an expression for the partition function of N ideal bosons in the explicit form. For this purpose we write the density matrix R_N as follows:

$$R_N(\mathbf{r}_1, \dots, \mathbf{r}_N | \mathbf{r}_1, \dots, \mathbf{r}_N) = \frac{1}{N!} \frac{1}{\lambda^{3N}} \Delta_N, \quad (7)$$

where

$$\Delta_N = \begin{vmatrix} 1 & K_{12} & K_{13} & \dots & K_{1N} \\ K_{21} & 1 & K_{23} & \dots & K_{2N} \\ \vdots & & \ddots & & \\ K_{N1} & K_{N2} & K_{N3} & \dots & 1 \end{vmatrix}_+, \quad (8)$$

and

$$K_{ij} \equiv K(r_{ij}) = e^{-\pi r_{ij}^2/\lambda^2}, \quad r_{ij} = |\mathbf{r}_{ij}|, \quad \mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j. \quad (9)$$

The subscript “+” means that every “−” in the expression for the determinant should be substituted with “+”. The partition function for free particles is

$$Z_N = \frac{1}{N!} \frac{1}{\lambda^{3N}} \int d\mathbf{r}_1 \dots \int d\mathbf{r}_N \Delta_N. \quad (10)$$

The “determinant” (8) might be written as [15]

$$\Delta_N = \Delta_{N-1}^{(1)} + \sum_{1 < i \leq N} K_{1i} K_{i1} \Delta_{N-2}^{(1,j)} + \sum_{1 < i < j \leq N} (K_{1i} K_{ij} K_{j1} + K_{1j} K_{ji} K_{i1}) \Delta_{N-3}^{(1,i,j)} + \dots, \quad (11)$$

The superscripts in Δ_{N-l} mean the lack of the correspondent rows and columns in the expression for Δ_{N-l} . Integrating (10) we obtain

$$Z_N = \frac{1}{N} \left\{ \frac{V}{\lambda^3} Z_{N-1} + \frac{1}{(\lambda^3)^2} Z_{N-2} \int d\mathbf{r}_1 \int d\mathbf{r}_2 K_{12} K_{21} + \frac{1}{(\lambda^3)^3} Z_{N-3} \int d\mathbf{r}_1 \int d\mathbf{r}_2 \int d\mathbf{r}_3 K_{12} K_{23} K_{31} + \dots \right\}, \quad (12)$$

After the Fourier transformation

$$K(r) = \frac{1}{V} \sum_{\mathbf{q}} e^{i\mathbf{q}\mathbf{r}} K_{\mathbf{q}}, \quad (13)$$

$$K_{\mathbf{q}} = \int d\mathbf{r} e^{-i\mathbf{q}\mathbf{r}} K(r) = \lambda^3 e^{-\lambda^2 q^2/4\pi} \quad (14)$$

the partition function is

$$Z_N = \frac{1}{\rho \lambda^3} \sum_{l=1}^N \frac{Z_{N-l}}{l^{3/2}}, \quad Z_0 \equiv 1. \quad (15)$$

One can use this expression to calculate the partition function of the system. We will make such calculations in the next section.

IV. MAIN RESULTS FOR THE FREE PARTICLES

It is possible to calculate heat capacities C_N for $N = 1, 2, 3, \dots$ particles sequentially using expressions (6) and (15). In addition, we can show explicitly the low- and high-temperature asymptotics for C_N , namely:

$$\left. \frac{C_N}{N} \right|_{T \rightarrow \infty} = \frac{3}{2}, \quad (16)$$

$$\left. \frac{C_N}{N} \right|_{T \rightarrow 0} = \frac{3}{2} \frac{1}{N} + A_N T^{3/2}, \quad A_N = \text{const with respect to } T. \quad (17)$$

In the case of $N = \infty$ this temperature dependencies coincide with the known results for the ideal Bose-gas [16]. For low temperatures one has

$$\left. \frac{C_N}{N} \right|_{N=\infty} = \frac{15}{4} \frac{\zeta(5/2)}{\zeta(3/2)} \left(\frac{T}{T_c} \right)^{3/2}; \quad (18)$$

the high-temperature limit being 3/2. The critical temperature T_c (the non-analyticity point on the heat capacity curve) in this case is defined from the equation

$$\rho \lambda_c^3 = \zeta(3/2), \quad (19)$$

where λ_c is the thermal de Broglie wavelength for T_c , and ζ is the Rieman ζ -function, $\zeta(3/2) = 2.612\dots$. Taking $\rho = 0.02185 \text{ rA}^{-3}$ for liquid ^4He we obtain $T_c = 3.138 \text{ K}$.

In this work we consider the systems with different N from 2 to 1200. As was mentioned above it is impossible to obtain the non-analyticity in the heat capacity, so we take the point of C_N maximum as the critical temperature $T_c^{(N)}$.

It appears that those values could be described by the linear dependence

$$\rho \left[\lambda_c^{(N)} \right]^3 \simeq 2.61 - \frac{c}{N^{1/3}}, \quad c \simeq 1.9 \quad (20)$$

with $\lambda_c^{(N)}$ being thermal de Broglie wavelength for $T_c^{(N)}$. One can easily notice that 2.61 is very close to $\zeta(3/2) = 2.612\dots$.

We also have a similiar correlation for the C_N maxima $C_N^{\max}/N$. So, one should expect that the limit $N \rightarrow \infty$ gives the correct curve (see




FIG. 1. Heat capacities in the ideal case.




FIG. 2. Temperature of the C_N/N maxima in the ideal case, Solid line shows dependence (20), circles correspond to the data from Table I.

The results are also given in Table I.

N	$\rho [\lambda_c^{(N)}]^3$	$C_N^{\max}/N$	N	$\rho [\lambda_c^{(N)}]^3$	$C_N^{\max}/N$
1	—	1.500	90	2.185	1.747
2	1.131	1.532	100	2.200	1.753
3	1.297	1.553	200	2.286	1.783
5	1.490	1.582	300	2.328	1.799
7	1.606	1.602	400	2.354	1.809
10	1.717	1.624	500	2.373	1.816
15	1.830	1.649	600	2.387	1.822
20	1.902	1.666	700	2.399	1.827
30	1.993	1.690	800	2.408	1.831
40	2.050	1.706	900	2.416	1.834
50	2.091	1.718	1000	2.423	1.837
60	2.123	1.727	1100	2.429	1.839
70	2.147	1.735	1200	2.434	1.842
80	2.168	1.742	∞	2.612	1.926

TABLE I. Main results for the ideal systems

As one can see from Fig. 1, good qualitative and satisfactory quantitative agreement is reached at $N = 100$. Although, typical elements of the heat capacity curve appear at smaller N , e. g., a clearly defined maximum is found at $N = 10$ –15. As we shall show further, these very values might be considered as the limit between “individual” and statistic behaviour of the system with N particles.

V. INTERACTION IN THE HARD-SPHERES POTENTIAL APPROXIMATION

In a many-body system, it is a very complicated problem to take into account the interaction and it has not as yet been completely solved. Specifically, the systems with a small number of particles give a possibility to consider this problem from the first principles, i. e., to include the interaction into the calculations explicitly. From the mathematical point of view, the computations become heavily complicated while the number of particles increases, and the direct interaction account soon turns out to be impossible.

We will proceed from the expression for the density matrix R_N in the free particles case. In the systems like ^4He small interatomic distances are essential when atoms behave almost as hard spheres. Thus, the wavefunction should vanish rapidly as soon as the distance between any two atoms approaches the hard sphere diameter.

Due to this we shall take the interaction into account by the substitution: of Δ_N in (10) with $\Delta_N P_N$ where P_N is a function of the particles coordinates:

$$Z_N = \frac{1}{N!} \frac{1}{\lambda^{3N}} \int d\mathbf{r}_1 \dots \int d\mathbf{r}_N \Delta_N P_N. \quad (21)$$

For the sake of simplicity we suppose the interaction to have a pair character since the obtained results are expected to be rather qualitative than quantitative. In the case of hard spheres potential we have

$$\begin{aligned}
P_N(\mathbf{r}_1, \dots, \mathbf{r}_N) &= \prod_{i=1}^{N-1} \prod_{j=2}^N P(i, j), \\
P(i, j) &= P(|\mathbf{r}_i - \mathbf{r}_j|) = P(r_{ij}), \\
P(i, j) &= 1 + f(i, j) = \begin{cases} 0, & r_{ij} \leq a \\ 1, & r_{ij} > a \end{cases},
\end{aligned} \quad (22)$$

a is the hard sphere diameter.

One can write the partition function as

$$Z_N = \frac{1}{\rho\lambda^3} \sum_{l=1}^N b'_l(N) Z_{N-l}, \quad Z_0 \equiv 1, \quad (23)$$

$$\lambda = \left(\frac{2\pi\beta\hbar^2}{m^*} \right)^{1/2}, \quad (24)$$

where m^* is the effective mass of ^4He atom, $m^* \simeq 1.7m$ [17].

We integrate (21) in two steps. The first one is *the circular approximation (CA)* and the second one is *the free-volume approximation (FVA)*.

1) CA is to the effect that only the interparticle interactions in the circle 1—2—3—...— N —1 are taken into account. It means we neglect the interaction between the first and the third, fourth, etc. particles, between the second and fourth, fifth, etc. particles, This makes it possible to integrate (21) using Fourier transformation as we did in (10). Using also FVA we obtain for the coefficients $b'_l(N)$:

$$b'_l(N) = \begin{cases} \left[1 - \frac{v_0}{V}(N-l)\right]^2 \left[1 - \frac{v_0}{V}(N-3)\right] \prod_{k=3}^{l-1} \left[1 - \frac{v_0}{V}(N-k)\right] I_{l-1}, & l > 3 \\ \left[1 - \frac{v_0}{V}(N-l)\right]^l I_{l-1}, & l \leq 3 \end{cases} \quad (25)$$

Here we have used such designations:

$$v_0 = \frac{4}{3}\pi a^3, \quad (26)$$

$$I_n = \begin{cases} 8(4\pi)^{n-1} \int_0^\infty dQ Q^2 \left(\frac{1}{4\pi} e^{-Q^2/4\pi} - \frac{1}{Q} \int_0^{a/\lambda} x \sin Qx e^{-\pi x^2} dx \right)^{n+1}, & n > 0 \\ 1, & n = 0 \end{cases} \quad (27)$$

2) In FVA we indirectly take into account the interactions neglected in CA. For this purpose each integral over $\mathbf{r}_i$, $i = 1, \dots, l$ is multiplied by the factor $(1 - nv_0/V)$ where n equals to the number of multipliers $P(i, j)$ which are not taken into account in the Fourier transformation. To clarify the above statement we consider the most simple non-trivial example with $N = 4$. According to (25) we have:

- $b'_1(4) = \left[1 - \frac{3v_0}{V}\right] I_0$. Integral over $\mathbf{r}_1$ “hooks” the following *three* multipliers: $P(1, 2)$, $P(1, 3)$, $P(1, 4)$.
- $b'_2(4) = \left[1 - \frac{2v_0}{V}\right]^2 I_1$. Integral over $\mathbf{r}_1$ “hooks” $P(1, 2)$, $P(1, 3)$, $P(1, 4)$ but $P(1, 2)$ is included into I_1 . Integral over $\mathbf{r}_2$ “hooks” more $P(2, 3)$ and $P(2, 4)$.
- $b'_3(4) = \left[1 - \frac{v_0}{V}\right]^3 I_2$. Each integral over $\mathbf{r}_i$ “hooks” additional $P(i, 2)$.
- $b'_4(4) = \left[1 - \frac{v_0}{V}\right]^2 I_3$. $P(1, 3)$ and $P(2, 4)$ are not included into I_3 .

Thus, FVA improves the circular approximation. But it is not possible to use it for large l and N because in this case a big number of $P(i, j)$ multipliers will not be integrated in the proper way but via formal factors.

In the calculations we assume the hard sphere diameter $a = 2.1$ rÅ. The heat capacity curves exhibit a fast approaching of the maximum point to a value from the region of 2.2–2.3 while N is increasing. The exact λ -transition temperature is 2.17 K.

As has already been mentioned, in the ideal case the linear (over $N^{-\varepsilon}$) dependence fits the sequence of C_N/N maxima quite well

$$\frac{C_N^{\max}}{N} \simeq C_\infty^{\max} - aN^{-\varepsilon}, \quad \varepsilon = 1/3. \quad (28)$$

Assuming the same form for the hard spheres system one can find $C_{\infty}^{\max} = 13.6$ and $\varepsilon = 0.0608$ making the maxima sequence as follows:

$$\frac{C_N^{\max}}{N} \simeq 13.6 - 12.9 N^{-\varepsilon}, \quad \varepsilon = 0.0608. \quad (29)$$

Although small N values of 12, 13 do not allow one to identify the behaviour at $N \rightarrow \infty$ we obtain the values of heat capacity maximum from the range of experimental data [18,19,7].




FIG. 3. Heat capacities in the free volume and circular approximation. The circles show the NIST C_V data [18].

One can write the exact expression for the partition function similar to that in the case of CA and FVA:

$$Z_N = \frac{1}{\rho \lambda^3} \sum_{l=1}^N b_l^*(N) Z_{N-l}, \quad Z_0 \equiv 1. \quad (30)$$

From expansion (11) and formula (21) we obtain (without approximations):

$$b_l^*(N) = \frac{1}{V} \frac{1}{(N-l)!} \frac{1}{(\lambda^3)^{N-1}} \int d\mathbf{r}_1 \dots \int d\mathbf{r}_l K_{12} \dots K_{l-1,l} K_{l1} \\ \times \prod_{i=1}^{l-1} \prod_{j=i+1}^l P(i,j) \mathcal{B}_{lN}(1, \dots, l), \quad (31)$$

$$\mathcal{B}_{lN}(1, \dots, l) = \int d\mathbf{r}_{l+1} \dots \int d\mathbf{r}_N \frac{\Delta_{N-l}^{(1, \dots, l)} P_{N-l}^{(1, \dots, l)} \prod_{i=1}^l \prod_{j=l+1}^N P(i,j)}{Z_{N-l}}, \quad (32)$$

$$P_{N-l}^{(1, \dots, l)} = \prod_{i=l+1}^{N-1} \prod_{j=i+1}^N P(i,j). \quad (33)$$

We also introduce the following designations to simplify further records:

$$R(1, 2) = \int d\mathbf{r}_3 P(1, 3)P(2, 3), \quad (34)$$

$$\tilde{K}_{ij} = K_{ij}P(i, j). \quad (35)$$

One can write the coefficients $b_i^*(N)$ independently of the interatomic potential, i. e. for any P_N :

$$b_1^*(1) = 1, \quad (36)$$

$$b_1^*(2) = \frac{1}{V^2} \int d\mathbf{r}_1 \int d\mathbf{r}_2 P(1, 2), \quad (37)$$

$$b_1^*(3) = \frac{\int d\mathbf{r}_1 \int d\mathbf{r}_2 P(1, 2)R(1, 2) (1 + K_{12}K_{21})}{\int d\mathbf{r}_1 \int d\mathbf{r}_2 P(1, 2) (1 + K_{12}K_{21})}, \quad (38)$$

$$b_2^*(2) = \frac{1}{V} \frac{1}{\lambda^3} \int d\mathbf{r}_1 \int d\mathbf{r}_2 K_{12}K_{21}P(1, 2), \quad (39)$$

$$b_2^*(3) = \frac{1}{V^2} \frac{1}{\lambda^3} \int d\mathbf{r}_1 \int d\mathbf{r}_2 K_{12}K_{21}P(1, 2)R(1, 2), \quad (40)$$

$$b_3^*(3) = \frac{1}{V} \frac{1}{(\lambda^3)^2} \int d\mathbf{r}_1 \int d\mathbf{r}_2 \int d\mathbf{r}_3 \tilde{K}_{12}\tilde{K}_{23}\tilde{K}_{31}. \quad (41)$$

These expressions might be written via one-dimensional integrals:

$$b_1^*(2) = \frac{4\pi}{V} \int_0^\infty P(r_{12})r_{12}^2 dr_{12}, \quad (42)$$

$$b_2^*(2) = \frac{4\pi}{\lambda^3} \int_0^\infty K_{12}^2 P(r_{12})r_{12}^2 dr_{12}, \quad (43)$$

$$R(r_{12}) = \frac{8}{r_{12}} \int_0^\infty \frac{dq}{q} \sin qr_{12} \left\{ \int_0^\infty r P(r) \sin qr dr \right\}^2, \quad (44)$$

$$b_3^*(3) = \frac{32\pi}{(\lambda^3)^2} \int_0^\infty \frac{dq}{q} \left\{ \int_0^\infty K_{12}P(r_{12})r_{12} \sin qr_{12} dr_{12} \right\}^3, \quad (45)$$

$$b_2^*(3) = \frac{4\pi}{V\lambda^3} \int_0^\infty K_{12}^2 P(r_{12})R(r_{12})r_{12}^2 dr_{12}. \quad (46)$$

In the hard spheres approximation integration is made simply. We obtain

$$b_1^*(1) = 1, \quad (47)$$

$$b_1^*(2) = 1 - \frac{v_0}{V}, \quad (48)$$

$$b_2^*(2) = \frac{1}{2^{3/2}} \left\{ 1 - \operatorname{erf} \left(\frac{a}{\lambda} \sqrt{2\pi} \right) \right\} + \frac{a}{\lambda} e^{-2\pi a^2/\lambda^2}, \quad (49)$$

$$b_1^*(3) = \frac{V - 3v_0 + \frac{81}{32} \frac{v_0^2}{V} + \lambda^3 b_2^*(3)}{V b_1^*(2) + \lambda^3 b_2^*(2)}, \quad (50)$$

$$b_2^*(3) = \frac{V - v_0}{V} \left\{ \frac{a}{\lambda} \left(e^{-2\pi a^2/\lambda^2} - 2e^{-8\pi a^2/\lambda^2} \right) + \frac{1}{2^{3/2}} \left(\operatorname{erf} \left(2\frac{a}{\lambda} \sqrt{2\pi} \right) - \operatorname{erf} \left(\frac{a}{\lambda} \sqrt{2\pi} \right) \right) \right\} \quad (51)$$

$$+ \frac{\pi a^2 \lambda}{V} \left\{ \frac{a^2}{\lambda^2} \left(e^{-2\pi a^2/\lambda^2} - 4e^{-8\pi a^2/\lambda^2} \right) + \frac{1}{2\pi} \left(e^{-2\pi a^2/\lambda^2} - e^{-8\pi a^2/\lambda^2} \right) \right\} \quad (52)$$

$$+ \frac{\pi \lambda^3}{12V} \left\{ e^{-2\pi a^2/\lambda^2} \left(\frac{a^4}{\lambda^4} + \frac{1}{\pi} \frac{a^2}{\lambda^2} + \frac{1}{2\pi^2} \right) - e^{-8\pi a^2/\lambda^2} \left(16 \frac{a^4}{\lambda^4} + \frac{4}{\pi} \frac{a^2}{\lambda^2} + \frac{1}{2\pi^2} \right) \right\} \quad (53)$$

$$+ \frac{V - 2v_0}{V} \left\{ \frac{1}{2^{3/2}} \left(1 - \operatorname{erf} \left(2\frac{a}{\lambda} \sqrt{2\pi} \right) \right) + 2\frac{a}{\lambda} e^{-8\pi a^2/\lambda^2} \right\}, \quad (54)$$

$$b_3^*(3) = 32\pi \int_0^\infty \frac{dQ}{Q} \left\{ \frac{Q}{4\pi} e^{-Q^2/4\pi} - \int_0^{a/\lambda} x \sin Qx e^{-\pi x^2} dx \right\}^3. \quad (55)$$

Our R coincides with the correspondent function from [20]:

$$R(r_{12}) = \int d\mathbf{r}_3 (1 + f_{13})(1 + f_{23}) = V - 2v_0 + g_1(r_{12}), \quad (56)$$

The function $g_1(r_{12})$ is given by

$$g_1(r) = \int d\mathbf{r}_3 f_{13}f_{23} = \begin{cases} \frac{2}{3}\pi a^3 \left(2 - \frac{3}{2}\frac{r}{a} + \frac{1}{8}\left(\frac{r}{a}\right)^3\right), & r < 2a \\ 0, & r \geq 2a \end{cases} \quad (57)$$

Meanwhile we cannot write exact expressions for greater N . The comparison of the exact results and those in CA and FVA is presented in Table II.

N	Approximation		Exact values	
	$T_c^{(N)}$	$C_N^{\max}/N$	$T_c^{(N)}$	$C_N^{\max}/N$
1	—	1.50	—	1.50
2	2.86	1.67	2.86	1.67
3	2.54	1.78	2.19	1.83
4	2.28	1.90	—	—
5	2.25	2.01	—	—
6	2.24	2.11	—	—
7	2.23	2.20	—	—
8	2.23	2.29	—	—
9	2.23	2.37	—	—
10	2.23	2.44	—	—
11	2.23	2.50	—	—
12	2.23	2.56	—	—
13*	2.23	2.61	—	—
18	2.23	2.82	—	—
∞	2.2–2.3	13.6**	—	—

* We suppose that the system of 12–13 particles behaves almost as a bulk since the atoms in ^4He are packed compactly enough to say that one atom is surrounded by 12 atoms. Therefore, 13 atoms constitute the first closed sphere (see Fig. 5).

** The experimental value is $\simeq 16$. The heat capacity in the λ -point was believed to be infinite for a long time but recent experiments show the finite peak [7].

TABLE II. Results for the interacting systems




FIG. 4. $C_N^{\max}/N$ values for the hard spheres system. Solid line shows dependence (29), circles correspond to the data from Table II.




FIG. 5. Helium-4 atoms as hard spheres. (a) The first closed sphere of 12 atoms surrounding 1 central atom. One atom is not seen since it is situated exactly behind the central atom. (b) Solid line shows the “surface of atom”, dashed line shows the “surface of alien atoms”, dashed-dotted line shows “surface of inaccessible space”. Thus, assuming the volume of “inaccessible space” $4\pi R^3/3$ to be the volume occupied by 1 atom one can obtain the hard sphere diameter $a \simeq 2.2$ rA.

Our expressions for the partition function have the correct form $Z_N^{(\text{cl})}$ in the classical limit ($\hbar \rightarrow 0$) [21].

$$Z_N^{(\text{cl})} = \frac{1}{N!} \frac{1}{\lambda^{3N/2}} \int d\mathbf{r}_1 \dots \int d\mathbf{r}_N \prod_{1 \leq i < j \leq N} P(i, j). \quad (58)$$

It means that the proposed method is applicable for a wide temperature range

CONCLUSION

It is shown that proposed method leads to a good agreement between the calculated results and the experimental data. We obtained not only qualitative but even satisfactory quantitative fit as well.

The expansions over $1/N$ obtained in this work demonstrate the existence of some parameter ε having the value of $1/3$ for the ideal system. We estimated $\varepsilon = 0.0608$ from the numerical analysis of the results. The theoretical calculation of this parameter will be our next problem along this line of research.

ACKNOWLEDGEMENTS

The authors appreciate the help of Dr. Daniel G. Friend with the helium-4 thermophysical data.

-
- [1] N. N. Bogoliubov, J. Phys. (USSR) **9**, 23 (1947).
 - [2] R. Brout, Phys. Rev. **131**, 899 (1963).
 - [3] R. P. Feynman, *Statistical Mechanics* (W. A. Benjamin, Reading, Mass., 1972).
 - [4] R. P. Feynman, Phys. Rev. **91**, 1291, 1301 (1953).

- [5] I. S. Langer, Phys. Rev., **167**, 183 (1968).
- [6] I. A. Vakarchuk, Teor. Mat. Fiz. **35**, 76 (1978); Teor. Mat. Fiz. **36**, 122 (1978) (in Russian).
- [7] J. A. Lipa, D. R. Swanson, J. A. Nissen, Phys. Rev. Lett. **76**, 944 (1996).
- [8] H. Kleinert, Phys. Rev. D **60**, 085001 (1999).
- [9] H. Kleinert, preprint cond-mat/9906107.
- [10] M. Campostrini, A. Pelissetto, P. Rossi, E. Vicari, Phys. Rev. B **61**, 5905 (2000).
- [11] D. M. Ceperley, Rev. Mod. Phys. **67**, 279 (1995).
- [12] Wenji Deng and P. M. Hui, Solid State Communications **104**, 797 (1997).
- [13] H.-J. Schmidt and J. Schnack, Physica A **260**, 479 (1998).
- [14] P. Borrmann, G. Franke, J. Chem. Phys. **98**, 2484 (1993).
- [15] I. O. Vakarchuk, J. Phys. Stud. (Lviv) **1**, 25 (1996); J. Phys. Stud. (Lviv) **1**, 156 (1997).
- [16] K. Huang, *Statistical Mechanics* (Wiley, New-York, 1987).
- [17] I. O. Vakarchuk, Visn. Lviv. un-tu, Ser. fiz. **26**, 29 (1993) (in Ukrainian).
- [18] V. D. Arp, R. D. McCarty, D. G. Friend, Natl. Inst. Stand. Technol., Tech. Note 1334 (revised) (1998).
- [19] M. Coleman and J. A. Lipa, Phys. Rev. Lett. **74**, 286 (1995).
- [20] B. R. A. Nijboer and L. Van Hove, Phys. Rev. **85**, 777 (1952).
- [21] A. Isihara, *Statistical Physics* (Academic Press, New York, 1971).